

# An Algebraic Criterion for Global Exponential Stability of Cohen-Grossberg Neural Networks with Time-varying Delays

Xinyuan Liang

College of Computer  
Science

Chongqing Technology and  
Business University  
Chongqing , P R China  
lxy@swsc.com.cn

Tian Wang

College of Computer  
Science

Chongqing Technology and  
Business University  
Chongqing , P R China  
ztewt@sohu.com

Zhengxia Wang

School of Science  
Chongqing Jiaotong  
University

Chongqing , P R China  
zxwang@163.com

Haixia Wu

Department of Computer  
and Modern Education  
Technology

Chongqing Education  
College  
Chongqing , P R China  
ammewu@163.com

**Abstract**—In this paper, by constructing an appropriate Lyapunov functional, sufficient criteria independent of the delays for global exponential stability of the network are derived. The algebra criteria are applicable for other neural network models. This results are less conservative and restrictive than previously known results and can be easily verified. And the result has overcome the obvious drawback that previous works neglect the signs of the connecting weights, and thus, do not distinguish the differences between excitatory and inhibitory connections. It is believed that the results are significant and useful for the design and applications of the Cohen-Grossberg model.

**Keywords**—Novel Criterion, Global Exponential Stability, Cohen-Grossberg Neural Network, Time-varying Delay

## I. INTRODUCTION

A large class of neural networks, which can function as stable content addressable memories or CAMs, are proposed by Cohen and Grossberg [1] and [2]. These Cohen-Grossberg networks were designed to include additive neural networks [3], [4], [7], [9], [10], [11], [16], [17], [21], [22], [23] and [24], later studied by Hopfield [3] and [4], and shunting neural networks. In general, the Cohen-Grossberg neural network model is described by the set of ordinary differential equations:

$$\dot{x}_i(t) = -a_i(x_i(t)) \left[ b_i(x_i(t)) - \sum_{j=1}^n w_{ij} f_j(x_j(t)) + J_i \right] \quad (1)$$

where  $i = 1, 2, \dots, n$ ,  $J_i$  denote the constant inputs from outside of the system and  $w_{ij}$  represent the connection weights.  $a_i(x_i)$  and  $b_i(x_i)$ , are the amplification functions and the self-signal functions, respectively, while  $f_j(x_j)$ ,  $j = 1, 2, \dots, n$ , are the activation functions. In the original analysis, Cohen and Grossberg assumed that the weight matrix  $W = (w_{ij})_{n \times n}$  is symmetric and the amplification functions  $a_i(x_i) > 0$ . Meanwhile, the activation functions  $f_j$  are assumed to be continuous, differentiable, monotonically increasing and bounded, such as the sigmoid-type function. For simplicity, the explanation of the lowercase  $i$  bound is omitted in the following equations.

In reality, time delays inevitably exist in biological and artificial neural networks due to the finite signal switching and transmission speed in a network. It is also important to incorporate time delay into the model equations of the network such as delayed cellular neural network, which can be used to solve problems like the processing of moving images [14] and [15]. Discrete delays were introduced into system (1) by considering the following system [5] and [8]:

$$\dot{x}_i(t) = -a_i(x_i(t)) \left[ b_i(x_i(t)) - \sum_{k=0}^K \sum_{j=1}^n v_{ij}^{(k)} f_j(x_j(t - \tau_k)) + J_i \right] \quad (2)$$

where the matrix  $V_k = (v_{ij}^{(k)})_{n \times n}$  represents the interconnections associated with delay  $\tau_k$  of which  $\tau_k$ ,  $k = 0, 1, \dots, K$ , are arranged such that  $0 = \tau_0 < \tau_1 < \dots < \tau_K$ . Furthermore, their global limit property needs to satisfy the requirements that the connection should possess certain amount of symmetry and the discrete delays are sufficiently small. Hence, the work of Ye et al. [8] cannot tell what would happen when the delays increase. We have mentioned above that large delay could destroy the stability of the equilibrium in a network. Even if the delay does not change the stability, it could affect the basin of attraction of the stable equilibrium. For this topic, interested readers may refer to the recent work [18], [29], [30] and [31]. Recently, for the delayed Hopfield networks [7], [9], [10] and [11], cellular neural networks [14] and [15] as well as BAM networks [12] and [13], some delay-independent criteria for the global asymptotic stability are established without assuming the monotonicity and the differentiability of the activation functions, nor the symmetry of the connections, [7] and [9]. Wang and Zou [5][6] studied the following system

$$\dot{x}_i(t) = -a_i(x_i(t)) \left[ b_i(x_i(t)) - \sum_{j=1}^n v_{ij} f_j(x_j(t - \tau_{ij})) + J_i \right] \quad (3)$$

where  $v_{ij} \geq 0$  are delays caused during the switching and transmission processes. Some criteria for the exponential stability of a unique equilibrium are obtained.

Liao [19] studied a modified Cohen-Grossberg model with discrete delays described by the differential difference equation of the form

$$\dot{x}_i(t) = -a_i(x_i(t)) \left[ b_i(x_i(t)) - \sum_{j=1}^n w_{ij} f_j(x_j(t)) - \sum_{j=1}^n v_{ij} f_j(x_j(t - \tau_{ij})) + J_i \right] \quad (4)$$

In this paper, we consider a modified Cohen-Grossberg model with multiple time-varying delays described by the differential difference equation of the form

$$\dot{x}_i(t) = -a_i(x_i(t)) \left[ b_i(x_i(t)) - \sum_{j=1}^n w_{ij} f_j(x_j(t)) - \sum_{k=0}^K \sum_{j=1}^n v_{ij}^{(k)} f_j(x_j(t - \tau_{ij}^{(k)}(t))) + J_i \right] \quad (5)$$

Unlike most of the previous authors, we did not confine to the symmetric connections. Moreover, we did not assume the monotonicity and differentiability of the activation functions. In this paper, the amplification functions only require to be continuous and positive. Moreover, the self-signal functions are not assumed to be differentiable. Thus a much broader connection topology of the network is allowed. The global exponential stability criteria of the Cohen-Grossberg neural network with time-varying delays were also obtained by constructing appropriate Lyapunov functions. And the result has overcome the obvious drawback that it neglects the signs of the connecting weights, and doesn't distinguish the differences between excitatory and inhibitory connections.

The rest of this paper is organized as follows. In Section 2, some preliminary analyses are given. By constructing a new Lyapunov functional, we studied the Cohen-Grossberg models with time-varying delays and gained some global exponential stability criteria in Section 3. Some comparisons and an example are given in Section 4 to demonstrate our main results. Finally, conclusions are drawn in Section 5.

## II. SOME PRELIMINARIES

Let  $R$  denotes the set of real numbers and if  $x \in R^n$ , then  $x^T = (x_1, x_2, \dots, x_n)$  denotes the transpose of  $x$ . Let  $R^{n \times n}$  denotes the set of  $n \times n$  real matrices. The initial conditions associated with (5) are in the form:

$$x_i(s) = \phi_i(s) \in C([- \tau, 0], R), s \in [- \tau, 0], i = 1, 2, \dots, n,$$

$$\tau = \max_{1 \leq i, j \leq n, 0 \leq k \leq K} \{ \tau_{ij}^{(k)}(t) \}.$$

For functions  $a_i(x_i)$  and  $b_i(x_i)$ ,  $i = 1, 2, \dots, n$ , the following assumptions are used.

(H<sub>1</sub>)  $a_i(u)$  are continuous and there exist positive constants  $\underline{a}_i$  and  $\bar{a}_i$  such that  $0 < \underline{a}_i \leq a_i(u) \leq \bar{a}_i$ , for all  $u \in R$ ,  $i = 1, 2, \dots, n$ .

(H<sub>2</sub>)  $b_i$  and  $b_i^{-1}$ ,  $i = 1, 2, \dots, n$ , are locally Lipschitz continuous and there exist  $\gamma_i > 0$ ,  $i = 1, 2, \dots, n$ , such that

$$u[b_i(u+x) - b_i(x)] \geq \gamma_i u^2$$

**Remark 1.** In [5], it is necessary that  $b_i'(x) > 0$ . However, our condition (H<sub>2</sub>) does not require that  $b_i(x)$ ,  $i = 1, 2, \dots, n$ , to

be continuous and differentiable. Moreover, our results are less conservative and restrictive than those given in [5].

Usually, the assumed activation functions are continuous, differentiable, and monotonically increasing and bounded, such as the sigmoid-type functions. However, as pointed out in [10], [11], [12], [13], [14], [15], [16] and [17], for certain purposes, non-monotonic and not necessarily smooth functions might be better candidates for the neuron activation functions in designing and implementing an artificial neural network. Note that in many electronic circuits, amplifiers possess neither monotonically increasing nor continuously differentiable input-output functions are frequently adopted.

Moreover, we assume that the activation functions  $f_i$ ,  $i = 1, 2, \dots, n$ , satisfy either (H<sub>3</sub>) or (H'<sub>3</sub>):

(H<sub>3</sub>) (i)  $f_i$ ,  $i = 1, 2, \dots, n$ , are bounded in  $R$ ;

(ii)  $0 < \frac{f_i(x+u) - f_i(x)}{u} \leq L_i$ ,  $i = 1, 2, \dots, n$ .

(H'<sub>3</sub>) (i)  $f_i$ ,  $i = 1, 2, \dots, n$ , are bounded in  $R$ ;

(ii)  $f_i(x+u) - f_i(x) \leq L_i |u|$ ,  $i = 1, 2, \dots, n$ .

(H<sub>4</sub>)  $\tau_{ij}^{(k)}(t) : [0, +\infty] \rightarrow [0, +\infty]$  is continuous, differentiable, and  $0 \leq \tau_{ij}^{(k)}(t) \leq \bar{\tau}$ ,  $\tau_{ij}^{(k)}(t) \leq q < 1$ ;

Hence, the hypothesis (H<sub>4</sub>) ensures that  $t - \tau_{ij}^{(k)}(t)$  has differential inverse function denoted by  $\phi_{ij}^{(k)}(t)$  and  $\phi_{ij}^{(k)}(t) > 0$ .

**Remark 2.** Note that unlike the requirement stated in [8],  $V = \sum_{k=0}^K V_k$  is not required to be symmetric in this paper. This means that our results are applicable to networks with a much broader connection structure.

Before stating the main results, we first need the following definitions and lemmas. For convenience, we introduce some notations. For a symmetric matrix,  $A > 0$  ( $< 0$ ) means that  $A$  is positive definite (negative definite).

**Definition 1.** For any continuous function  $h: R \rightarrow R$ , Dini's time-derivative of  $h(t)$  is defined as

$$D^+ h(t) = \limsup_{\theta \rightarrow 0^+} \frac{h(t+\theta) - h(t)}{\theta}.$$

It is easy to see that if  $h(t)$  is locally Lipschitz, then  $D^+ h(t) < \infty$ .

**Definition 2.** If there exist  $k > 0$  and  $\gamma(k) > 1$ , such that

$$\|x(t)\| \leq \gamma(k) e^{-kt} \sup_{-\tau \leq \theta \leq 0} \|x(\theta)\|, \forall t > 0,$$

system (5) is considered as exponential stable, and  $k$  is called the degree of exponential stability.

We note that  $x^*$  is an equilibrium of system (5) if and only if  $x^* = [x_1^*, x_2^*, \dots, x_n^*]^T \in R$  is a solution of the following equations

$$b_i(x_i^*) - \sum_{j=1}^n w_{ij} f_j(x_j^*) - \sum_{k=0}^K \sum_{j=1}^n v_{ij}^{(k)} f_j(x_j^*) + J_i \quad (6)$$

Similar to the discussion in [5], we can easily obtain the following results.

**Lemma 1.** *If  $(H_1)$ - $(H_2)$  and  $(H_3)$  or  $(H'_3)$  hold, then for every input  $J$ , there exists an equilibrium point for system (5).*

Let  $x^*$  be an equilibrium of system (5) and  $z(t) = x(t) - x^*$ . Substituting  $x(t) = z(t) + x^*$  into system (5) leads to

$$\begin{aligned} \dot{z}_i(t) = & -a_i(z_i(t) + x_i^*) \left[ b_i(z_i(t) + x_i^*) - \sum_{j=1}^n w_{ij} f_j(z_j(t) + x_j^*) \right. \\ & \left. - \sum_{k=0}^K \sum_{j=1}^n v_{ij}^{(k)} f_j(z_j(t - \tau_{ij}^{(k)}(t)) + x_j^*) + J_i \right] \end{aligned} \quad (7)$$

By (7), system (5) can be rewritten as

$$\dot{z}_i(t) = -\alpha_i(z_i(t)) \left[ \beta_i(z_i(t)) - \sum_{j=1}^n w_{ij} g_j(z_j(t)) - \sum_{k=0}^K \sum_{j=1}^n v_{ij}^{(k)} g_j(z_j(t - \tau_{ij}^{(k)}(t))) \right] \quad (8)$$

where

$$\begin{aligned} \alpha_i(z_i(t)) &= a_i(z_i(t) + x_i^*) \\ \beta_i(z_i(t)) &= b_i(z_i(t) + x_i^*) - b_i(x_i^*) \\ g_i(z_i(t)) &= f_i(z_i(t) + x_i^*) - f_i(x_i^*) \end{aligned} \quad (9)$$

It is obvious that  $x^*$  is global exponential stable for system (5) if and only if the trivial solution  $z = 0$  of system (7) is global exponential stable. Moreover, the uniqueness of the equilibrium of (5) follows from its global exponential stability.

From assumption  $H'_3$ , we have

$$(H_5) \quad |g_i(z_i)| \leq L_i |z_i|, \quad g_i(0) = 0, \quad i = 1, 2, \dots, n.$$

Then, it is easy to see that

$$z_i g_i(z_i) \leq L_i z_i^2.$$

**Definition 3.** For any scalar  $a \in \mathbb{R}$ ,  $a^+$  is defined as

$$a^+ = \max\{0, a\}. \quad (10)$$

or

$$a^+ = |a|. \quad (11)$$

**Lemma 2.** Any scalar  $a \in \mathbb{R}$ , there exist scalar  $a^+$ , we hold the inequality  $a \leq a^+$ .

**Proof.** Any scalar  $a \in \mathbb{R}$ , we have

$$a \leq \max\{0, a\}, \text{ and } a \leq |a|.$$

It is easy to see that  $a \leq a^+$ .

### III. MAIN RESULTS

**Theorem 1.** *Consider the delayed system (5) and assume that conditions  $(H_1)$ - $(H_4)$  are satisfied. If there exist  $\psi_{ij}, \rho_{ij}, \zeta_{ij}, \eta_{ij} \in [0, 1]$ , and positive constants  $r_1, r_2, d_i, i, j = 1, 2, \dots, n$ , and  $\sigma > 0$ , and the following condition holds:*

$$\begin{aligned} \Delta = & \max_{1 \leq i \leq n} \left\{ \left[ d_i \left( \gamma_i - \frac{\sigma}{\alpha_i} \right) \right]^{-1} \left[ \sum_{j=1}^n \left( r_1 d_i (w_{ij}^+)^{2\psi_{ij}} L_j^{2\rho_{ij}} + \frac{1}{r_1} d_j \right. \right. \right. \\ & \left. \left. (w_{ji}^+)^{2(1-\psi_{ji})} L_i^{2(1-\rho_{ji})} \right) + \sum_{k=0}^K \sum_{j=1}^n \left( r_2 d_i ((v_{ij}^{(k)})^+)^{2\zeta_{ij}} L_j^{2\eta_{ij}} (1-q)^{-1} \right. \right. \\ & \left. \left. e^{2\sigma v_{ij}^{(k)}(t)} + \frac{1}{r_2} d_j ((v_{ji}^{(k)})^+)^{2(1-\zeta_{ji})} L_i^{2(1-\eta_{ji})} \right) \right\} < 2 \end{aligned} \quad (12)$$

then, for every input  $J$ , the equilibrium  $x^*$  for system (5) is global exponential stable. This implies that there exists constants  $\eta \geq 1$  such that every solution of (5) satisfies

$$\sum_{i=1}^n (x_i(t) - x_i^*)^2 \leq \eta e^{-2\sigma t} \sup_{s \in [-\tau, 0]} (\phi_i(s) - x_i^*)^2, \quad t > 0 \quad (13)$$

where

$$\begin{aligned} \eta \leq & \max_{1 \leq i \leq n} \left\{ \frac{d_i}{\alpha_i} \left( 1 + \frac{\alpha_i}{2\sigma} (1 - e^{-2\sigma\tau}) \sum_{k=0}^K \sum_{j=1}^n \frac{1}{r_2} ((v_{ij}^{(k)})^+)^{2(1-\zeta_{ij})} L_j^{2(1-\eta_{ij})} \right) \right\} \\ & \left/ \min_{1 \leq i \leq n} \left( \frac{d_i}{\alpha_i} \right) \right. \end{aligned}$$

**Proof.** We construct the following nonnegative functional as the Lyapunov functional candidate:

$$\begin{aligned} V(z(t)) = & \sum_{i=1}^n d_i \left\{ e^{2\sigma t} \int_0^{\tau_i(t)} \frac{s}{\alpha_i(s)} ds + \right. \\ & \left. 2 \sum_{k=0}^K \sum_{j=1}^n \frac{1}{r_2} ((v_{ij}^{(k)})^+)^{2(1-\zeta_{ij})} L_j^{2(1-\eta_{ij})} \int_{t-\tau_{ij}^{(k)}(t)}^t z_j^2(s) e^{2\sigma s} ds \right\} \end{aligned} \quad (14)$$

By computing its the Dini's time-derivative of  $V(z(t))$  along the trajectories of (8) and making use of the Cauchy inequality (i.e,  $ra^2 + b^2/r \geq 2ab, r > 0$ ), we have

$$\begin{aligned} D^+ V(z(t)) = & \sum_{i=1}^n d_i \left\{ 2e^{2\sigma t} \frac{z_i(t)}{\alpha_i(z_i(t))} \dot{z}_i(t) + 4\sigma e^{2\sigma t} \int_0^{\tau_i(t)} \frac{s}{\alpha_i(s)} ds + \right. \\ & \sum_{k=0}^K \sum_{j=1}^n \frac{1}{r_2} ((v_{ij}^{(k)})^+)^{2(1-\zeta_{ij})} L_j^{2(1-\eta_{ij})} z_j^2(t) e^{2\sigma t} \\ & \left. - \sum_{k=0}^K \sum_{j=1}^n \frac{1}{r_2} ((v_{ij}^{(k)})^+)^{2(1-\zeta_{ij})} L_j^{2(1-\eta_{ij})} (1 - \dot{\tau}_{ij}^{(k)}(t)) z_j^2(t - \tau_{ij}^{(k)}(t)) e^{2\sigma(t - \tau_{ij}^{(k)}(t))} \right\} \\ = & \sum_{i=1}^n d_i \left\{ -2e^{2\sigma t} z_i(t) \left[ \beta_i(z_i(t)) - \sum_{j=1}^n w_{ij} g_j(z_j(t)) - \sum_{k=0}^K \sum_{j=1}^n v_{ij}^{(k)} g_j(z_j(t - \tau_{ij}^{(k)}(t))) \right] \right. \\ & \left. + 4\sigma e^{2\sigma t} \int_0^{\tau_i(t)} \frac{s}{\alpha_i(s)} ds + \sum_{k=0}^K \sum_{j=1}^n \frac{1}{r_2} ((v_{ij}^{(k)})^+)^{2(1-\zeta_{ij})} L_j^{2(1-\eta_{ij})} z_j^2(t) e^{2\sigma t} \right\} \end{aligned}$$

$$\begin{aligned}
& - \sum_{k=0}^K \sum_{j=1}^n \frac{1}{r_2} ((v_{ij}^{(k)})^+)^{2(1-\zeta_{ij})} L_j^{2(1-\eta_j)} (1 - \tau_{ij}^{(k)}(t)) z_j^2(t - \tau_{ij}^{(k)}(t)) e^{2\sigma(t - \tau_{ij}^{(k)}(t))} \Big\} \\
& \leq \sum_{i=1}^n d_i e^{2\sigma} \left\{ -2z_i(t) \beta_i(z_i(t)) + 2 \sum_{j=1}^n w_{ij} z_i(t) g_j(z_j(t)) + 2 \sum_{k=0}^K \sum_{j=1}^n v_{ij}^{(k)} z_i(t) \right. \\
& \quad \left. g_j(z_j(t - \tau_{ij}^{(k)}(t))) + \frac{2\sigma}{\alpha_i} z_i^2(t) + \sum_{k=0}^K \sum_{j=1}^n \frac{1}{r_2} ((v_{ij}^{(k)})^+)^{2(1-\zeta_{ij})} L_j^{2(1-\eta_j)} z_j^2(t) - \right. \\
& \quad \left. \sum_{k=0}^K \sum_{j=1}^n \frac{1}{r_2} ((v_{ij}^{(k)})^+)^{2(1-\zeta_{ij})} L_j^{2(1-\eta_j)} (1-q) z_j^2(t - \tau_{ij}^{(k)}(t)) e^{-2\sigma \tau_{ij}^{(k)}(t)} \right\} \\
& \leq \sum_{i=1}^n d_i e^{2\sigma} \left\{ -2\gamma_i z_i^2(t) + 2 \sum_{j=1}^n w_{ij}^+ L_j |z_i(t)| |z_j(t)| + 2 \sum_{k=0}^K \sum_{j=1}^n (v_{ij}^{(k)})^+ L_j |z_i(t)| \right. \\
& \quad \left. |z_j(t - \tau_{ij}^{(k)}(t))| + \frac{2\sigma}{\alpha_i} z_i^2(t) + \sum_{k=0}^K \sum_{j=1}^n \frac{1}{r_2} ((v_{ij}^{(k)})^+)^{2(1-\zeta_{ij})} L_j^{2(1-\eta_j)} z_j^2(t) \right. \\
& \quad \left. - \sum_{k=0}^K \sum_{j=1}^n \frac{1}{r_2} ((v_{ij}^{(k)})^+)^{2(1-\zeta_{ij})} L_j^{2(1-\eta_j)} (1-q) z_j^2(t - \tau_{ij}^{(k)}(t)) e^{-2\sigma \tau_{ij}^{(k)}(t)} \right\} \\
& = \sum_{i=1}^n d_i e^{2\sigma} \left\{ -2 \left( \gamma_i - \frac{\sigma}{\alpha_i} \right) z_i^2(t) + 2 \sum_{j=1}^n ((w_{ij}^+)^{w_{ij}} L_j^{\rho_{ij}} |z_i(t)|) ((w_{ij}^+)^{1-w_{ij}} L_j^{1-\rho_{ij}} |z_j(t)|) + \right. \\
& \quad \left. 2 \sum_{k=0}^K \sum_{j=1}^n \left[ ((v_{ij}^{(k)})^+)^{\zeta_{ij}} L_j^{\eta_{ij}} (1-q)^{-\frac{1}{2}} e^{\sigma \tau_{ij}^{(k)}(t)} |z_i(t)| \right] \right. \\
& \quad \left. \left[ (v_{ij}^{(k)})^+ \right]^{-\zeta_{ij}} L_j^{1-\eta_{ij}} (1-q)^{\frac{1}{2}} |z_j(t - \tau_{ij}^{(k)}(t))| e^{-\sigma \tau_{ij}^{(k)}(t)} \right] + \right. \\
& \quad \left. \sum_{k=0}^K \sum_{j=1}^n \frac{1}{r_2} ((v_{ij}^{(k)})^+)^{2(1-\zeta_{ij})} L_j^{2(1-\eta_j)} z_j^2(t) - \right. \\
& \quad \left. \sum_{k=0}^K \sum_{j=1}^n \frac{1}{r_2} ((v_{ij}^{(k)})^+)^{2(1-\zeta_{ij})} L_j^{2(1-\eta_j)} (1-q) z_j^2(t - \tau_{ij}^{(k)}(t)) e^{-2\sigma \tau_{ij}^{(k)}(t)} \right\} \\
& \leq \sum_{i=1}^n d_i e^{2\sigma} \left\{ -2 \left( \gamma_i - \frac{\sigma}{\alpha_i} \right) z_i^2(t) + \sum_{j=1}^n r_1 (w_{ij}^+)^{2w_{ij}} L_j^{2\rho_{ij}} z_i^2(t) + \right. \\
& \quad \sum_{j=1}^n \frac{1}{r_1} (w_{ij}^+)^{2(1-w_{ij})} L_j^{2(1-\rho_{ij})} z_j^2(t) + \sum_{k=0}^K \sum_{j=1}^n r_2 ((v_{ij}^{(k)})^+)^{2\zeta_{ij}} L_j^{2\eta_{ij}} (1-q)^{-1} e^{2\sigma \tau_{ij}^{(k)}(t)} z_i^2(t) \\
& \quad + \sum_{k=0}^K \sum_{j=1}^n \frac{1}{r_2} ((v_{ij}^{(k)})^+)^{2(1-\zeta_{ij})} L_j^{2(1-\eta_j)} (1-q) z_j^2(t - \tau_{ij}^{(k)}(t)) e^{-2\sigma \tau_{ij}^{(k)}(t)} \\
& \quad + \sum_{k=0}^K \sum_{j=1}^n \frac{1}{r_2} ((v_{ij}^{(k)})^+)^{2(1-\zeta_{ij})} L_j^{2(1-\eta_j)} z_j^2(t) - \\
& \quad \left. \sum_{k=0}^K \sum_{j=1}^n \frac{1}{r_2} ((v_{ij}^{(k)})^+)^{2(1-\zeta_{ij})} L_j^{2(1-\eta_j)} (1-q) z_j^2(t - \tau_{ij}^{(k)}(t)) e^{-2\sigma \tau_{ij}^{(k)}(t)} \right\} \\
& = \sum_{i=1}^n d_i \left( \gamma_i - \frac{\sigma}{\alpha_i} \right) e^{2\sigma} \left\{ -2 + \left[ d_i \left( \gamma_i - \frac{\sigma}{\alpha_i} \right) \right]^{-1} \left[ \sum_{j=1}^n (r_1 d_i (w_{ij}^+)^{2w_{ij}} L_j^{2\rho_{ij}} \right. \right. \right. \\
& \quad \left. \left. \frac{1}{r_1} d_j (w_{ji}^+)^{2(1-w_{ji})} L_i^{2(1-\rho_{ji})} \right) + \sum_{k=0}^K \sum_{j=1}^n \left( r_2 d_i ((v_{ij}^{(k)})^+)^{2\zeta_{ij}} L_j^{2\eta_{ij}} (1-q)^{-1} e^{2\sigma \tau_{ij}^{(k)}(t)} \right. \right. \\
& \quad \left. \left. + \frac{1}{r_2} d_j ((v_{ji}^{(k)})^+)^{2(1-\zeta_{ji})} L_i^{2(1-\eta_{ji})} \right) \right] \Big\} z_i^2(t) \\
& = \max_{1 \leq i \leq n} \left\{ d_i \left( \gamma_i - \frac{\sigma}{\alpha_i} \right) (-2 + \Delta) \right\} \sum_{i=1}^n z_i^2(t) e^{2\sigma} = -B \sum_{i=1}^n z_i^2(t) e^{2\sigma} < 0 \quad (15)
\end{aligned}$$

This implies that  $V(z(t)) \leq V(z(0))$  for  $t > 0$ , and

$$\begin{aligned}
V(z(0)) & = \sum_{i=1}^n d_i \left\{ 2 \int_0^{\tau_i(0)} \frac{s}{\alpha_i(s)} ds + \sum_{k=0}^K \sum_{j=1}^n \frac{1}{r_2} ((v_{ij}^{(k)})^+)^{2(1-\zeta_{ij})} \right. \\
& \quad \left. L_j^{2(1-\eta_j)} \int_{-\tau_{ij}^{(k)}(0)}^0 z_j^2(s) e^{2\sigma s} ds \right\} \\
& \leq \sum_{i=1}^n d_i \left\{ \frac{1}{\alpha_i} z_i^2(0) + \sum_{k=0}^K \sum_{j=1}^n \frac{1}{r_2} ((v_{ij}^{(k)})^+)^{2(1-\zeta_{ij})} L_j^{2(1-\eta_j)} \right. \\
& \quad \left. \sup_{s \in [-\tau, 0]} \left\{ z_j^2(s) \right\} \int_{-\tau_{ij}^{(k)}(0)}^0 e^{2\sigma s} ds \right\} \\
& \leq \max_{1 \leq i \leq n} \left\{ \frac{d_i}{\alpha_i} \left( 1 + \frac{\alpha_i}{2\sigma} (1 - e^{-2\sigma \tau}) \right) \right. \\
& \quad \left. \sum_{k=0}^K \sum_{j=1}^n \frac{1}{r_2} ((v_{ij}^{(k)})^+)^{2(1-\zeta_{ij})} L_j^{2(1-\eta_j)} \right\} \sum_{i=1}^n \sup_{s \in [-\tau, 0]} (\phi_i(s) - x_i^*)^2
\end{aligned}$$

By (14), we can easily obtain

$$V(z(0)) \geq e^{2\sigma} \sum_{i=1}^n 2d_i \int_0^{\tau_i(t)} \frac{s}{\alpha_i(s)} ds \geq \min_{1 \leq i \leq n} \left( \frac{d_i}{\alpha_i} \right) e^{2\sigma} \sum_{i=1}^n z_i^2(t) \quad (16)$$

Hence, we have the following results

$$\sum_{i=1}^n (x_i(t) - x_i^*)^2 \leq \eta e^{-2\sigma} \sum_{i=1}^n \sup_{s \in [-\tau, 0]} (\phi_i(s) - x_i^*)^2, t > 0 \quad (17)$$

Now, by a standard Lyapunov-type theorem for functional differential equation, the trivial solution of (8) with (9) is global exponential stable for  $\Delta < 2$ . Therefore,  $x^*$  is global exponential stable for system (5). This completes the proof of Theorem 1.

**Remark 3.** In [19][26][27], the author presented a sufficient condition for global exponential stability of (2) and (4). However, compared with Theorem 1 above the previous result has an obvious drawback: it neglects the signs of the connecting weights, and thus, does not distinguish the differences between excitatory ( $w_{ij} > 0$ ) and inhibitory ( $w_{ij} < 0$ ) connections [25].

If we set  $K=0$ , in Theorem 1, then the following corollary is immediate.

**Corollary 1.** Consider the delayed system (5) and assume that conditions  $(H_1)$ - $(H_4)$  are satisfied. If there exist  $\psi_{ij}, \rho_{ij}, \zeta_{ij}, \eta_{ij} \in [0, 1]$ , and positive constants  $r_1, r_2, d_i, i, j = 1, 2, \dots, n$ , and  $\sigma > 0$ , and the following condition holds:

$$\begin{aligned}
\Delta & = \max_{1 \leq i \leq n} \left\{ \left[ d_i \left( \gamma_i - \frac{\sigma}{\alpha_i} \right) \right]^{-1} \left[ \sum_{j=1}^n \left( r_1 d_i (w_{ij}^+)^{2w_{ij}} L_j^{2\rho_{ij}} + \frac{1}{r_1} d_j \right. \right. \right. \\
& \quad \left. \left. (w_{ji}^+)^{2(1-w_{ji})} L_i^{2(1-\rho_{ji})} \right) + \sum_{j=1}^n \left( r_2 d_i ((v_{ij}^+)^+)^{2\zeta_{ij}} L_j^{2\eta_{ij}} (1-q)^{-1} \right. \right. \\
& \quad \left. \left. e^{2\sigma \tau_{ij}^{(k)}(t)} + \frac{1}{r_2} d_j ((v_{ji}^+)^+)^{2(1-\zeta_{ji})} L_i^{2(1-\eta_{ji})} \right) \right] \Big\} < 2 \quad (18)
\end{aligned}$$

then, for every input  $J$ , the equilibrium  $x^*$  for system (5) is global exponential stable.

If we set  $w_{ij} = 0$ ,  $i, j = 1, 2, \dots, n$ , in Theorem 1, then the following corollary is immediate.

**Corollary 2.** Consider the delayed system (2) and (5) with pure delays and assume that conditions (H<sub>1</sub>)-(H<sub>4</sub>) are satisfied. If there exist  $\psi_{ij}, \rho_{ij}, \zeta_{ij}, \eta_{ij} \in [0, 1]$ , and positive constants  $r_1, r_2, d_i, i, j = 1, 2, \dots, n$ , and  $\sigma > 0$ , and the following condition holds:

$$\Delta = \max_{1 \leq i \leq n} \left\{ \left[ d_i \left( \gamma_i - \frac{\sigma}{\alpha_i} \right) \right]^{-1} \left[ \sum_{k=0}^K \sum_{j=1}^n \left( r_2 d_i ((v_{ij}^{(k)})^+)^{2\zeta_{ij}} L_j^{2\eta_{ij}} \right. \right. \right. \\ \left. \left. \left. (1-q)^{-1} e^{2\sigma\tau_{ij}^{(k)}(t)} + \frac{1}{r_2} d_j ((v_{ji}^{(k)})^+)^{2(1-\zeta_{ji})} L_i^{2(1-\eta_{ji})} \right) \right] \right\} < 2 \quad (19)$$

then, for every input  $J$ , the equilibrium  $x^*$  for system (2) and (5) is global exponential stable.

When we set  $a_i(x) = 1$ , the model becomes the following neural networks with multiple time-varying delays:

$$\dot{x}_i(t) = -b_i(x_i(t)) + \sum_{j=1}^n w_{ij} f_j(x_j(t)) + \sum_{k=0}^K \sum_{j=1}^n v_{ij}^{(k)} f_j(x_j(t - \tau_{ij}^{(k)})) + J_i \quad (20)$$

When we set  $a_i(x) = 1$ ,  $b_i(x) = \gamma_i x$  in system (5), where  $\gamma_i > 0$ ,  $i = 1, 2, \dots, n$ , the model becomes the following cellular neural networks with multiple time-varying delays:

$$\dot{x}_i(t) = -\gamma_i x_i(t) + \sum_{j=1}^n w_{ij} f_j(x_j(t)) + \sum_{k=0}^K \sum_{j=1}^n v_{ij}^{(k)} f_j(x_j(t - \tau_{ij}^{(k)})) + J_i \quad (21)$$

Hence, by virtue of Theorem 1, we can immediately obtain the following result:

**Corollary 3.** Consider the delayed system (20) and (21) and assume that conditions (H<sub>1</sub>)-(H<sub>4</sub>) are satisfied. If there exist  $\psi_{ij}, \rho_{ij}, \zeta_{ij}, \eta_{ij} \in [0, 1]$ , and positive constants  $r_1, r_2, d_i, i, j = 1, 2, \dots, n$ , and  $\sigma > 0$ , and the following condition holds:

$$\Delta = \max_{1 \leq i \leq n} \left\{ \left[ d_i (\gamma_i - \sigma) \right]^{-1} \left[ \sum_{j=1}^n \left( r_1 d_i (w_{ij}^+)^{2\psi_{ij}} L_j^{2\rho_{ij}} + \frac{1}{r_1} d_j \right. \right. \right. \\ \left. \left. \left. (w_{ji}^+)^{2(1-\psi_{ji})} L_i^{2(1-\rho_{ji})} \right) + \sum_{k=0}^K \sum_{j=1}^n \left( r_2 d_i ((v_{ij}^{(k)})^+)^{2\zeta_{ij}} L_j^{2\eta_{ij}} (1-q)^{-1} \right. \right. \right. \\ \left. \left. \left. e^{2\sigma\tau_{ij}^{(k)}(t)} + \frac{1}{r_2} d_j ((v_{ji}^{(k)})^+)^{2(1-\zeta_{ji})} L_i^{2(1-\eta_{ji})} \right) \right] \right\} < 2 \quad (22)$$

then, for every input  $J$ , the equilibrium  $x^*$  for system (20) and (21) is global exponential stable.

#### IV. REMARKS AND AN EXAMPLE

In this Section, we will give an example to illustrate that the conditions given in this paper are less conservative and restrictive as than previously known results.

Example1 Consider the following system:

$$\begin{pmatrix} \dot{x}_1(t) \\ \dot{x}_2(t) \end{pmatrix} = - \begin{pmatrix} 4 + \sin x_1 & 0 \\ 0 & 2 + \cos x_2 \end{pmatrix} \begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix} \begin{pmatrix} x_1(t) \\ x_2(t) \end{pmatrix} \\ - \begin{pmatrix} -1 & 0.25 \\ 0.5 & -0.85 \end{pmatrix} \begin{pmatrix} \tanh(x_1(t - \tau_1)) \\ \tanh(2x_2(t - \tau_2)) \end{pmatrix} + \begin{pmatrix} J_1 \\ J_2 \end{pmatrix} \quad (23)$$

In this example,  $L_1=1, L_2=2, \gamma_1=2, \gamma_2=2, \bar{\alpha}_1=5, \alpha_1=3, \bar{\alpha}_2=3, \alpha_2=1$ . By virtue of condition in [5], it is easy to compute that

$$\theta_1 = \frac{\alpha_1}{\bar{\alpha}_1} \gamma_1 - L_1 \sum_{k=0}^1 \sum_{j=1}^2 |v_{j1}^{(k)}| = -0.3, \quad \text{and} \\ \theta_2 = \frac{\alpha_2}{\bar{\alpha}_2} \gamma_2 - L_2 \sum_{k=0}^1 \sum_{j=1}^2 |v_{j2}^{(k)}| = -1.533$$

Therefore, we have  $\theta = \min\{\theta_1, \theta_2\} = -1.533 < 0$ , hence, the conditions in [5] are not satisfied.

We can easily calculate

$$\theta_1 = \frac{1}{\gamma_1} \sum_{k=0}^1 \sum_{j=1}^2 (L_j |v_{1j}^{(k)}| + L_1 |v_{j1}^{(k)}|) = 1.5, \quad \text{and} \\ \theta_2 = \frac{1}{\gamma_2} \sum_{k=0}^1 \sum_{j=1}^2 (L_j |v_{2j}^{(k)}| + L_1 |v_{j2}^{(k)}|) = 2.2$$

Therefore, we have  $\theta = \max\{\theta_1, \theta_2\} = 2.2 > 2$ , hence, the conditions of Theorem 1 in [20] are not satisfied.

We can easily calculate

$$\theta_1 = \alpha_1 \gamma_1 - L_1 \sum_{k=0}^1 \sum_{j=1}^2 \alpha_j |v_{j1}^{(k)}| = 2.5, \quad \text{and} \\ \theta_2 = \alpha_2 \gamma_2 - L_2 \sum_{k=0}^1 \sum_{j=1}^2 \alpha_j |v_{j2}^{(k)}| = -1.2$$

Therefore, we have  $\theta = \min\{\theta_1, \theta_2\} = -1.2 < 0$ , hence, the conditions in [28] are not satisfied.




Figure 1. System (23) with initial state  $x(0) = [0.5, -0.3]^T$ .

However, if  $\psi_{ij} = \rho_{ij} = \zeta_{ij} = \eta_{ij} = 1/2$ ,  $w_{ij}^+ = \max\{w_{ij}, 0\}$ ,  $d_1=d_2=r_1=r_2=1$ ,  $\sigma = 0.4624$  in Corollary 3, then we can calculate  $\Delta_1=1.992, \Delta_2=1.145$ . Therefore, we have  $\Delta = \max\{\Delta_1, \Delta_2\} = 1.992 < 2$ , hence, the conditions given in Corollary 3 are hold. For numerical simulation, System (23) with initial states  $x(0) = [0.5, -0.3]^T$ , inputs  $J_1 = 1.5, J_2 = -1.5$ , and delay parameters  $\tau_1=1$  and  $\tau_2=2$  are considered. The existence of an unique equilibrium point,  $x^* = [-0.4568, 0.3732]^T$ , and the global

asymptotic stability of system (23) are guaranteed by means of computer simulations. Fig.1 depicted the time response of the state variables  $x_1(t)$  and  $x_2(t)$  for this case. Moreover, the results are independent of the delay parameters.

## V. CONCLUSIONS

In this paper, the criteria for the global exponential stability of a class of Cohen-Grossberg neural networks with multiple time-varying delays have been derived. The results have been shown to be the generalization and improvement of existing results reported recently in the literature for the cases with delays. Analyses have also shown that the neuronal input-output activation function and the self-signal function only need to satisfy, respectively, conditions  $(H_1)$ ,  $(H_2)$ ,  $(H_3)$  and  $(H_4)$  given in this paper, but do not need to be continuous, differentiable, monotonically increasing and bounded, as usually required by other analyzing methods. Novel stability conditions are stated in simple algebraic forms so that their verification and applications are straightforward and convenient. The criteria are independent of the magnitudes of the delays. We only require  $a_i(x)$  be continuous and positive. At the same time, we think over the signs of the entries of the connection matrices. Thus the differences between excitatory and inhibitory effects are considered. An example is given to demonstrate that our criteria are less conservative and restrictive than previously known results.

## ACKNOWLEDGMENTS

The work described here was partially supported by the Natural Science Foundation of China (Grant No. 60574024, 60573047, 60703035), the Post-doctoral Science Foundation of China ((Grant No.20060400714), the Natural Science Foundation Project of CQ CSTC (Grant No. 2007BB2130, 2006EA2011) and the Science and Technology Project of Chongqing Education Committee (Grant No. kj070708).

## REFERENCES

- [1] M.A. Cohen and S. Grossberg, Absolute stability and global pattern formation and parallel memory storage by competitive neural networks, *IEEE Trans Syst, Man Cybernetics*, SMC 13 (1981), pp. 815-821.
- [2] S. Grossberg, Nonlinear neural networks: principles, mechanisms, and architectures, *Neural Networks* 1 (1988), pp. 17-61.
- [3] J.J. Hopfield, Neuronal networks and physical systems with emergent collective computational abilities, *Proc National Acad Sci* 79 (1982), pp. 2554-2558.
- [4] J.J. Hopfield, Neurons with graded response have collective computational properties like those of two-state neurons, *Proc National Acad Sci* 81 (1984), pp. 3058-3092.
- [5] L. Wang and X. Zou, Harmless delays in Cohen-Grossberg neural networks, *Physica D* 170 (2002), pp. 162-173.
- [6] Wang and X. Zou, Exponential of Cohen-Grossberg neural networks, *Neural Networks* 15 (2002), pp. 415-422.
- [7] Z. Guan, G. Chen and Y. Qin, On equilibria, stability, and instability of Hopfield neural networks, *IEEE Trans Neural Networks* 11 (2000), pp. 534-540.
- [8] H. Ye, A.N. Michel and K. Wang, Qualitative analysis of Cohen-Grossberg neural networks with multiple delays, *Phys Rev E* 51 (1995), pp. 2611-2618.
- [9] M. Joy, Results concerning the absolute stability of delayed neural networks, *Neural Networks* 13 (2000), pp. 613-616.
- [10] X.F. Liao and J.B. Yu, Robust stability for interval Hopfield neural networks with time delays, *IEEE Trans Neural Networks* 9 (1998), pp. 1042-1046.
- [11] X.F. Liao, K.W. Wong, Z.F. Wu and G. Chen, Novel robust stability criteria for interval delayed Hopfield neural networks, *IEEE Trans CAS-I* 48 (2001), pp. 1355-1359.
- [12] X.F. Liao and J.B. Yu, Qualitative analysis of bi-directional associative memory networks with time delays, *Int J Circuit Theory Appl* 26 (1998), pp. 219-229.
- [13] X.F. Liao, J.B. Yu and G. Chen, Novel stability criteria for bi-directional associative memory neural networks with time delays, *Int J Circuit Theory Appl* 30 (2002), pp. 519-546.
- [14] X.F. Liao, K.W. Wong and J.B. Yu, Novel stability conditions for cellular neural networks with time delay, *Int J Bifur Chaos* 11 (2001), pp. 1853-1864.
- [15] X.F. Liao, Z.F. Wu and J.B. Yu, Stability analyses for cellular neural networks with continuous delay, *J Comput Appl Math* 143 (2002), pp. 29-47.
- [16] X.F. Liao, G. Chen and E.N. Sanchez, LMI-based approach for asymptotically stability analysis of delayed neural networks, *IEEE Trans CAS-I* 49 (2002), pp. 1033-1039.
- [17] X.F. Liao, G. Chen and E.N. Sanchez, Delay-dependent exponential stability analysis of delayed neural networks: an LMI approach, *Neural Networks* 15 (2002), pp. 855-866.
- [18] K. Pakdaman, C. Grotta-Ragazzo, C.P. Malta and J.F. Viber, Effect of delay on the boundary of the basin of attraction in a self-excited single neuron, *Neural Comput* 9 (1997), pp. 319-336.
- [19] X.F. Liao, C.D. Li and K.W. Wong, Criteria for exponential stability of Cohen-Grossberg neural networks, *Neural Networks* (2004), pp. 1401-1414.
- [20] Fenghua Tu, Xiaofeng Liao, Harmless delays for global asymptotic stability of Cohen-Grossberg neural networks, *Chaos, Solitons & Fractals* 26 (2005), pp. 927-933.
- [21] Qiang Zhang, Xiaopeng Wei, Jin Xu, Stability of delayed cellular neural networks. *Chaos Solitons & Fractals* 2007;31 :514-520.
- [22] Qiang Zhang, Xiaopeng Wei, Jin Xu, Delay-dependent exponential stability of cellular neural networks with time-varying delays. *Chaos Solitons & Fractals* 2005;23: 1363-1369.
- [23] Chuandong Li, Xiaofeng Liao, Rong Zhang, Delay-dependent exponential stability analysis of bi-directional associative memory neural networks with time delay: an LMI approach. *Chaos Solitons & Fractals* 2005;24: 1119-1134.
- [24] Chuandong Li, Xiaofeng Liao, Rong Zhang, A global exponential robust stability criterion for interval delayed neural networks with variable delays. *Neurocomputing* 2006;69: 803-809.
- [25] Vanden Driessche P, Wu J, Zou X. Stabilization role of inhibitory self-connections in a delayed neural network. *Physica D* 2001;150: 84-90.
- [26] Jinde Cao, Boundedness and stability for Cohen-Grossberg neural network with time-varying delays, *Journal of Mathematical Analysis and Application* 2004; 296: 665-685.
- [27] Sabri Arik, Zeynep Orman, Global stability analysis of Cohen-Grossberg neural networks with time varying delays, *Physics Letters A* 2005; 341: 410-421.
- [28] Jin Deng, Daoyi Xu, Zhichun Yang, DengJin(2004)-Global exponential stability of Cohen-Grossberg neural networks with multiple time-varying delays, the 11th IEEE International Conference on Electronics, Circuits and Systems, ICECS 2004; 2004, pp. 230-233
- [29] Wei Wu, BaoTong Cui, Min Huang, Global asymptotic stability of Cohen-Grossberg neural networks with constant and variable delays, *Chaos, Solitons & Fractals* 2007; 33: 1355-1361
- [30] Wei Wu, BaoTong Cui, Min Huang, Global asymptotic stability of delayed Cohen-Grossberg neural networks, *Chaos, Solitons & Fractals* 2007; 34: 872-877
- [31] Wei Wu, BaoTong Cui, XuYang Lou, Some criteria for asymptotic stability of Cohen-Grossberg neural networks with time-varying delay, *Neurocomputing* 2007; 70: 1085-1088