

On fuzzy preordered sets and monotone Galois connections

F. García-Pardo, I.P. Cabrera, P. Cordero, M. Ojeda-Aciego
 Universidad de Málaga. Andalucía Tech. Spain
 {fgarciap, ipcabrera, pcordero, aciego}@uma.es

Abstract—In this work, we focus on the study of necessary and sufficient conditions in order to ensure the existence (under some constraints) of monotone Galois connections between fuzzy preordered sets.

I. INTRODUCTION

Monotone Galois connections (also called adjunctions) between two mathematical structures provide a means of linking both theories allowing for mutual cooperative advantages. They, together with their antitone counterparts, have played an important role in computer science because its many applications, both theoretical and practical, and in mathematics because of its ability to link apparently very disparate worlds; this is why Denecke, Ern e, and Wismath stated in their monograph [14] that *Galois connections provide the structure-preserving passage between two worlds of our imagination*.

Finding a monotone Galois connection (or Galois connection) between two fields is extremely useful, since it provides a strong link between both theories allowing for mutual synergistic advantages. The algebraic study of complexity of valued constraints, for instance, has been studied in terms of establishing a Galois connection [10].

A number of results can be found in the literature concerning sufficient or necessary conditions for a Galois connection between ordered structures to exist. The main result of this paper is related to the existence and construction of the right adjoint to a given mapping f , but in a more general framework. It is worth to recall that, in a fuzzy setting, reflexivity and antisymmetry are conflicting properties [4] and, whereas some authors [20] opted for dropping reflexivity, our choice in this case has been to ignore antisymmetry and, therefore, consider fuzzy preorders.

Hence, our initial setting is to consider a mapping $f: A \rightarrow B$ from a fuzzy preordered set A into an unstructured set B , and then characterize those situations in which B can be fuzzy preordered and an isotone mapping $g: B \rightarrow A$ can be built such that the pair (f, g) is a monotone Galois connection.

In a previous work [25], we provided the set of necessary conditions for a monotone Galois connection to exist between fuzzy preordered sets. The main contribution in this paper is to prove that the necessary conditions are also sufficient.

The structure of this work is the following: in the next section, we introduce the preliminary definitions and results, essentially notions related to fuzzy preorderings and to Galois

connections, and some results which will be later needed. Section III introduces several lemmas which allow to simplify the presentation of the proof of the main result in Section IV, where the construction of the right adjoint is given based on the set of necessary conditions already known from [25].

II. PRELIMINARIES

As usual, as underlying structure for considering the generalization to a fuzzy framework, we will consider a *residuated lattice* $\mathbb{L} = (L, \vee, \wedge, \top, \perp, \otimes, \rightarrow)$, i.e. $(L, \vee, \wedge, 0, 1)$ is a bounded lattice, $(L, \otimes, 1)$ is a commutative monoid and (\otimes, \rightarrow) is an adjoint pair ($a \otimes b \leq c$ iff $a \leq b \rightarrow c$).

An \mathbb{L} -fuzzy set is a mapping from the universe set, say X , to the lattice L , i.e. $X: U \rightarrow L$, where $X(u)$ means the degree in which u belongs to X .

Given X and Y two \mathbb{L} -fuzzy sets, X is said to be included in Y , denoted as $X \subseteq Y$, if $X(u) \leq Y(u)$ for all $u \in U$.

An \mathbb{L} -fuzzy binary relation on U is an \mathbb{L} -fuzzy subset of $U \times U$, that is $\rho_U: U \times U \rightarrow L$, and it is said to be:

- Reflexive if, for each $a \in U$,

$$\rho_U(a, a) = \top$$

- Transitive if, for each $a, b, c \in U$,

$$\rho_U(a, b) \otimes \rho_U(b, c) \leq \rho_U(a, c)$$

- Symmetric if, for each $a, b \in U$,

$$\rho_U(a, b) = \rho_U(b, a)$$

- Antisymmetric if, for each $a, b \in U$,

$$\rho_U(a, b) = \rho_U(b, a) = \top \text{ implies } a = b$$

Definition 1 (Fuzzy poset / fuzzy preordered set):

- An \mathbb{L} -fuzzy poset is a pair $\mathbb{U} = (U, \rho_U)$ in which ρ_U is a reflexive, antisymmetric and transitive \mathbb{L} -fuzzy relation on U .
- An \mathbb{L} -fuzzy preordered set is a pair $\mathbb{U} = (U, \rho_U)$ in which ρ_U is a reflexive and transitive \mathbb{L} -fuzzy relation on U .
- A crisp (pre-)ordering can be given in U by $a \leq_U b$ if and only if $\rho_U(a, b) = \top$.

From now on, when no confusion arises, we will omit the prefix “ \mathbb{L} -”.

Definition 2: For every element $a \in U$, the extension to the fuzzy setting of the notions of *upset* and *downset* of the element a are defined by $a^\uparrow, a^\downarrow: U \rightarrow L$ where $a^\downarrow(u) = \rho_U(u, a)$ and $a^\uparrow(u) = \rho_U(a, u)$ for all $u \in U$.

An element $a \in U$ is an *upper bound* for a fuzzy set X if $X \subseteq a^\downarrow$. The (crisp) set of upper bounds of X is denoted by $UB(X)$. An element $a \in U$ is a *maximum* for a fuzzy set X if it is an upper bound and $X(a) = \top$.

The definitions of *lower bound* and *minimum* are similar. Note that, because of antisymmetry, maximum and minimum elements are necessarily unique.

Definition 3: Let $\mathbb{A} = (A, \rho_A)$ and $\mathbb{B} = (B, \rho_B)$ be fuzzy posets.

- 1) A mapping $f: A \rightarrow B$ is said to be *isotone* if $\rho_A(a_1, a_2) \leq \rho_B(f(a_1), f(a_2))$ for each $a_1, a_2 \in A$.
- 2) A mapping $f: A \rightarrow A$ is said to be *inflationary* if $\rho_A(a, f(a)) = \top$ for all $a \in A$.
- 3) A mapping $f: A \rightarrow A$ is *deflationary* if, for all $a \in A$, $\rho_A(f(a), a) = \top$.

Definition 4 (Monotone Galois connection): Let $\mathbb{A} = (A, \rho_A)$ and $\mathbb{B} = (B, \rho_B)$ be fuzzy posets, and two mappings $f: A \rightarrow B$ and $g: B \rightarrow A$. The pair (f, g) forms an *monotone Galois connection* between A and B , denoted $(f, g): \mathbb{A} \rightleftharpoons \mathbb{B}$ if, for all $a \in A$ and $b \in B$, the equality $\rho_A(a, g(b)) = \rho_B(f(a), b)$ holds.

Notation 1: From now on, we will use the following notation, for a mapping $f: A \rightarrow B$ and a fuzzy subset Y of B , the fuzzy set $f^{-1}(Y)$ is defined as $f^{-1}(Y)(a) = Y(f(a))$, for all $a \in A$.

Finally, we recall the following theorem which states different equivalent forms to define a monotone Galois connection between fuzzy posets.

Theorem 1 ([24]): Let $\mathbb{A} = (A, \rho_A)$, $\mathbb{B} = (B, \rho_B)$ be fuzzy posets, and two mappings $f: A \rightarrow B$ and $g: B \rightarrow A$. The following conditions are equivalent:

- 1) $(f, g): \mathbb{A} \rightleftharpoons \mathbb{B}$.
- 2) f and g are isotone, $g \circ f$ is inflationary, and $f \circ g$ is deflationary.
- 3) $f(a)^\uparrow = g^{-1}(a^\uparrow)$ for all $a \in A$.
- 4) $g(b)^\downarrow = f^{-1}(b^\downarrow)$ for all $b \in B$.
- 5) f is isotone and $g(b) = \max f^{-1}(b^\downarrow)$ for all $b \in B$.
- 6) g is isotone and $f(a) = \min g^{-1}(a^\uparrow)$ for each $a \in A$.

The next theorem characterizes the situation in which a mapping from a fuzzy poset to an unstructured set has a right adjoint (between fuzzy posets).

Theorem 2 ([26]): Let (A, ρ_A) be a fuzzy poset and a mapping $f: A \rightarrow B$. Let A_f be the quotient set over the kernel relation $a \equiv_f b$ iff $f(a) = f(b)$.

Then, there exists a fuzzy order ρ_B in B and a map $g: B \rightarrow A$ such that $(f, g): \mathbb{A} \rightleftharpoons \mathbb{B}$ if and only if the following conditions hold:

- 1) There exists $\max[a]_f$ for all $a \in A$.
- 2) $\rho_A(a_1, a_2) \leq \rho_A(\max[a_1]_f, \max[a_2]_f)$, for all $a_1, a_2 \in A$.

III. BUILDING MONOTONE GALOIS CONNECTIONS BETWEEN FUZZY PREORDERED SETS

In this section we start the generalization of Theorem 2 above to the framework of fuzzy preordered sets.

The construction will follow that given in [28] as much as possible. Therefore, we need to define a suitable fuzzy version of the p -kernel relation.

Firstly, we need to set the corresponding fuzzy notion of transitive closure of a fuzzy relation, and this is done via the definition below:

Definition 5 (Transitive closure): Given a fuzzy relation $S: U \times U \rightarrow L$, for all $n \in \mathbb{N}$, the iterations $S^n: U \times U \rightarrow L$ are recursively defined by the base case $S^1 = S$ and, then,

$$S^n(a, b) = \bigvee_{x \in U} \left(S^{n-1}(a, x) \otimes S(x, b) \right)$$

The *transitive closure* of S is a fuzzy relation $S^{tr}: U \times U \rightarrow L$ defined by

$$S^{tr}(a, b) = \bigvee_{n=1}^{\infty} S^n(a, b)$$

The relation \approx_A allows for getting rid of the absence of antisymmetry, by linking together elements which are ‘almost coincident’; formally, the relation \approx_A is defined on a fuzzy preordered set (A, ρ_A) as follows:

$$(a_1 \approx_A a_2) = \rho_A(a_1, a_2) \otimes \rho_A(a_2, a_1) \quad \text{for } a_1, a_2 \in A$$

The *kernel equivalence relation* \equiv_f associated to a mapping $f: A \rightarrow B$ is defined as follows for $a_1, a_2 \in A$:

$$(a_1 \equiv_f a_2) = \begin{cases} \perp & \text{if } f(a_1) \neq f(a_2) \\ \top & \text{if } f(a_1) = f(a_2) \end{cases}$$

Definition 6 (Fuzzy p -kernel): Let $\mathbb{A} = (A, \rho_A)$ be a fuzzy preordered set, and $f: A \rightarrow B$ a mapping. The *fuzzy p -kernel* relation \cong_A is the fuzzy equivalence relation obtained as the transitive closure of the union of the relations \approx_A and \equiv_f . Notice that the fuzzy equivalence classes $[a]_{\cong_A}: A \rightarrow L$ are fuzzy sets, whose definition is the following:

$$[a]_{\cong_A}(x) = (x \cong_A a)$$

Lemma 1: Let $\mathbb{A} = (A, \rho_A)$ be a fuzzy preordered set, and $f: A \rightarrow B$ a mapping. Then, $a_1 \cong_A a_2 = \top$ if and only if $[a_1]_{\cong_A} = [a_2]_{\cong_A}$.

PROOF: Consider $a_1, a_2 \in A$ such that $a_1 \cong_A a_2 = \top$, and let us prove that $[a_1]_{\cong_A}(u) = [a_2]_{\cong_A}(u)$ for all $u \in A$. Given $u \in A$, by using the neutral element of the product, and symmetry and transitivity of \cong_A , we have that

$$\begin{aligned} (a_1 \cong_A u) &= \top \otimes (a_1 \cong_A u) \\ &= (a_2 \cong_A a_1) \otimes (a_1 \cong_A u) \leq (a_2 \cong_A u) \end{aligned}$$

Similarly, $(a_2 \cong_A u) \leq (a_1 \cong_A u)$ and, therefore, $[a_1]_{\cong_A}(u) = [a_2]_{\cong_A}(u)$ for all $u \in A$. \square

All the preliminary notions about fuzzy posets introduced in the previous section carry over fuzzy preordered sets. Note, however, that there is an important difference which justifies the introduction of special terminology concerning maximum or minimum element of a fuzzy subset X : due to the absence

of antisymmetry, *there exists a crisp set of maxima* (resp. minima) for X , not necessarily a singleton, which we will denote $\text{p-max}(X)$ (resp., $\text{p-min}(X)$).

The following theorem states the different equivalent characterizations of the notion of monotone Galois connection between fuzzy preordered sets. As expected, the general structure of the definitions is preserved, but those concerning the actual definition of the adjoints have to be modified by using the notions of p-maximum and p-minimum.

Theorem 3 ([24]): Let $\mathbb{A} = (A, \rho_A)$ and $\mathbb{B} = (B, \rho_B)$ be two fuzzy preordered sets, and $f: \mathbb{A} \rightarrow \mathbb{B}$ and $g: \mathbb{B} \rightarrow \mathbb{A}$ be two mappings. The following statements are equivalent:

- 1) $(f, g): \mathbb{A} \rightleftharpoons \mathbb{B}$.
- 2) f and g are isotone, and $g \circ f$ is inflationary, $f \circ g$ is deflationary.
- 3) $f(a)^\uparrow = g^{-1}(a^\uparrow)$ for all $a \in A$.
- 4) $g(b)^\downarrow = f^{-1}(b^\downarrow)$ for all $b \in B$.
- 5) f is isotone and $g(b) \in \text{p-max } f^{-1}(b^\downarrow)$ for all $b \in B$.
- 6) g is isotone and $f(a) \in \text{p-min } g^{-1}(a^\uparrow)$ for all $a \in A$.

The following definitions recall the notion of Hoare ordering between crisp subsets, and then we introduce an alternative statement in the subsequent lemma:

Definition 7: Given a fuzzy preordered set (A, ρ_A) , and C, D crisp subsets of A , we define the following relations

- $(C \sqsubseteq_W D) = \bigvee_{c \in C} \bigvee_{d \in D} \rho_A(c, d)$
- $(C \sqsubseteq_H D) = \bigwedge_{c \in C} \bigvee_{d \in D} \rho_A(c, d)$
- $(C \sqsubseteq_S D) = \bigwedge_{c \in C} \bigwedge_{d \in D} \rho_A(c, d)$

Lemma 2 ([25]): Consider a fuzzy preordered set (A, ρ_A) , and $X, Y \subseteq A$ such that $\text{p-min } X \neq \emptyset \neq \text{p-min } Y$, then

$$\begin{aligned} (\text{p-min } X \sqsubseteq_W \text{p-min } Y) &= (\text{p-min } X \sqsubseteq_H \text{p-min } Y) \\ &= (\text{p-min } X \sqsubseteq_S \text{p-min } Y) \end{aligned}$$

and their value coincides with $\rho_A(x, y)$ for any $x \in \text{p-min } X$ and $y \in \text{p-min } Y$.

In [27], given a crisp poset (A, \leq_A) and a map $f: A \rightarrow B$, it was proved that there exists an ordering \leq_B in B and a map $g: B \rightarrow A$ such that (f, g) is a crisp monotone Galois connection between posets from (A, \leq_A) to (B, \leq_B) *if and only if*

- (I) There exists $\max([a]_{\equiv_f})$ for all $a \in A$.
- (II) $a_1 \leq_A a_2$ implies $\max([a_1]_{\equiv_f}) \leq_A \max([a_2]_{\equiv_f})$, for all $a_1, a_2 \in A$.

where \equiv_f is the kernel relation associated to f .

These two conditions are closely related to the different characterizations of the notion of monotone Galois connection, as stated in Theorem 1 (items 5 and 6); specifically, condition (I) above states that if $b \in B$ and $f(a) = b$, then necessarily $g(b) = \max([a]_{\equiv_f})$, whereas condition (II) is related to the isotonicity of both f and g .

Later, in [28], the previous result was extended to give necessary and sufficient conditions to ensure similar result in the framework of crisp preordered sets. Specifically, it was

proved that given any (crisp) preordered set $\mathbb{A} = (A, \lesssim_A)$ and a mapping $f: A \rightarrow B$, there exists a preorder $\mathbb{B} = (B, \lesssim_B)$ and $g: B \rightarrow A$ such that (f, g) forms a crisp monotone Galois connection between \mathbb{A} and \mathbb{B} *if and only if* there exists a subset S of A such that the following conditions hold:

- (i) $S \subseteq \bigcup_{a \in A} \text{p-max}[a]_{\cong_A}$
- (ii) $\text{p-min}(UB[a]_{\cong_A} \cap S) \neq \emptyset$, for all $a \in A$.
- (iii) For $a_1, a_2 \in A$, if $a_1 \lesssim_A a_2$ then

$$\left(\text{p-min}(UB[a_1]_{\cong_A} \cap S) \right) \sqsubseteq_H \left(\text{p-min}(UB[a_2]_{\cong_A} \cap S) \right)$$

It is worth to mention that in the conditions above all the notions used are the corresponding crisp versions of those defined in this paper.

In some sense, the conditions (i), (ii), (iii) reflect the considerations given in the previous paragraph, but the different underlying ordered structure leads to a different formalization. Formally, condition (I) above is split into (i) and (ii), since in a preordered setting, if $b \in B$ and $f(a) = b$, then $g(b)$ needs not be in the same class as a but being maximum in its class, as (i) states. However, the latter condition is too weak and (ii) provides exactly the remaining requirements needed in order to adequately reproduce the desired properties for g . Now, condition (iii) is just the rephrasing of (II) in terms of the properties described in (ii).

Finally, in [25], it was proved that the natural extension of the previous conditions to the fuzzy case are also necessary conditions to ensure the existence of a monotone Galois connection between fuzzy preordered sets. Specifically,

Theorem 4: Given fuzzy preordered sets $\mathbb{A} = (A, \rho_A)$ and $\mathbb{B} = (B, \rho_B)$, and mappings $f: A \rightarrow B$ and $g: B \rightarrow A$ such that $(f, g): \mathbb{A} \rightleftharpoons \mathbb{B}$ then

- 1) $gf(A) \subseteq \bigcup_{a \in A} \text{p-max}[a]_{\cong_A}$
- 2) $\text{p-min}(UB[a]_{\cong_A} \cap gf(A)) \neq \emptyset$, for all $a \in A$.
- 3) For all $a_1, a_2 \in A$, $\rho_A(a_1, a_2) \leq \left(\text{p-min}(UB[a_1]_{\cong_A} \cap gf(A)) \sqsubseteq_H \text{p-min}(UB[a_2]_{\cong_A} \cap gf(A)) \right)$.

As a consequence of the previous theorem, a necessary condition for f to be a left adjoint is the existence of a subset $S \subseteq A$ such that the following conditions hold for all $a, a_1, a_2 \in A$:

$$S \subseteq \bigcup_{a \in A} \text{p-max}[a]_{\cong_A} \quad (1)$$

$$\varphi_S(a) \neq \emptyset, \quad (2)$$

$$\rho_A(a_1, a_2) \leq (\varphi_S(a_1) \sqsubseteq_H \varphi_S(a_2)) \quad (3)$$

where

$$\varphi_S(a) \stackrel{\text{def}}{=} \text{p-min}(UB[a]_{\cong_A} \cap S). \quad (4)$$

Remark 1: Notice that, by Lemma 2, $(\varphi_S(a_1) \sqsubseteq_H \varphi_S(a_2)) = \rho_A(x, y)$ for any $x \in \varphi_S(a_1)$ and $y \in \varphi_S(a_2)$, and this justifies that, in order to simplify the notation, we write $\rho_A(\varphi_S(a_1), \varphi_S(a_2))$ instead of $(\varphi_S(a_1) \sqsubseteq_H \varphi_S(a_2))$. The main contribution in this paper is to show the converse, namely, that the conditions above are also sufficient so that f is a left adjoint.

IV. CONSTRUCTION OF THE RIGHT ADJOINT

In this section, given $f: A \rightarrow B$ with the conditions above, we will construct a fuzzy preordering on B together with a mapping $g: B \rightarrow A$, which will turn out to be a right adjoint to f .

Definition 8: Consider a fuzzy preordered set $\mathbb{A} = (A, \rho_A)$ together with a mapping $f: A \rightarrow B$ and a subset $S \subseteq A$ satisfying the ambient hypotheses (1), (2) and (3).

For all $a_0 \in A$, we define the fuzzy relation $\rho_B^{a_0}: B \times B \rightarrow L$ as follows

$$\rho_B^{a_0}(b_1, b_2) = \rho_A(\varphi_S(a_1), \varphi_S(a_2))$$

where $a_i \in f^{-1}(b_i)$ if $f^{-1}(b_i) \neq \emptyset$ and $a_i = a_0$ otherwise, for each $i \in \{1, 2\}$.

Notice that the definition might depend largely on the possible choices of a_i ; the following lemma, based on Remark 1, shows that the value of $\rho_B^{a_0}$ actually is independent of these choices.

Lemma 3: The fuzzy relation $\rho_B^{a_0}$ is well-defined, and it is a fuzzy preordering in B .

PROOF: The definition does not depend on the choice of preimages a_i since, if other preimages \bar{a}_i would have been chosen, then $(a_i \equiv_f \bar{a}_i) = \top$ and, hence, by Lemma 1, the fuzzy sets corresponding to the equivalence classes $[a_i]_{\cong_A}$ and $[\bar{a}_i]_{\cong_A}$ would coincide and $\varphi_S(a_i) = \varphi_S(\bar{a}_i)$. Moreover, by Remark 1, we have that

$$\rho_A(\varphi_S(a_1), \varphi_S(a_2)) = \rho_A(x, y)$$

for any $x \in \varphi(a_1)$ and $y \in \varphi(a_2)$, whose value is independent from the choice of x and y .

From the reflexivity of ρ_A , it is straightforward that $\rho_B^{a_0}$ is reflexive. Finally, it is just a matter of easy computations to check that $\rho_B^{a_0}$ is transitive. \square

We can now focus on the definition of suitable mappings $g: B \rightarrow A$ such that (f, g) forms an adjoint pair.

Lemma 4: Let $\mathbb{A} = (A, \rho_A)$ be a fuzzy preordered set, $f: A \rightarrow B$ be a mapping and S be a subset of A satisfying the ambient hypotheses (1), (2) and (3). Given $a_0 \in A$, then there exists a mapping $g: B \rightarrow A$ such that $(f, g): (A, \rho_A) \rightleftharpoons (B, \rho_B^{a_0})$ where $\rho_B^{a_0}$ is the fuzzy preordering introduced in Definition 8.

PROOF: There is a number of suitable definitions of $g: B \rightarrow A$, and all of them can be specified as follows:

- (C1) If $b \in f(A)$, then $g(b)$ is any element in $\varphi_S(x_b)$ for some $x_b \in f^{-1}(b)$.
- (C2) If $b \notin f(A)$, then $g(b)$ is any element in $\varphi_S(a_0)$.

The existence of g is clear by the axiom of choice, since for all $b \in f(A)$, the sets $f^{-1}(b)$ are nonempty (so x_b can be chosen for all $b \in f(A)$) and, moreover, by ambient hypothesis (2), $\varphi_S(x_b)$ and $\varphi_S(a_0)$ are nonempty as well.

Now, we have to prove that g is a right adjoint to f , that is, for all $a \in A$ and $b \in B$ the following equality holds

$$\rho_B^{a_0}(f(a), b) = \rho_A(a, g(b))$$

By definition of $\rho_B^{a_0}$ (see Definition 8), we have that

$$\rho_B^{a_0}(f(a), b) = \rho_A(\varphi_S(a), \varphi_S(w))$$

where w satisfies either $w \in f^{-1}(b)$ if $b \in f(A)$ (therefore, we can choose w to be x_b above) or, otherwise, $w = a_0$. In either case, $g(b) \in \varphi_S(w)$ by construction (namely, (C1) and (C2)). Thus,

$$\rho_B^{a_0}(f(a), b) = \rho_A(x, g(b)) \text{ for any } x \in \varphi_S(a) \quad (5)$$

The proof will be finished if we show that, fixing $x \in \varphi_S(a)$, we can show the equality $\rho_A(x, g(b)) = \rho_A(a, g(b))$.

Firstly, by definition of φ_S , see (4), note that $x \in \varphi_S(a)$ implies $\rho_A(a, x) = \top$ and, hence, we have that

$$\rho_A(x, g(b)) = \rho_A(a, x) \otimes \rho_A(x, g(b)) \leq \rho_A(a, g(b)) \quad (6)$$

For the other inequality, using ambient hypothesis (3), we have

$$\rho_A(a, g(b)) \leq \rho_A(\varphi_S(a), \varphi_S(g(b))) = \rho_A(x, y) \quad (7)$$

for any $x \in \varphi_S(a)$ and $y \in \varphi_S(g(b))$.

Since $y \in \varphi_S(g(b))$ we have that $\rho_A(y, \alpha) = \top$ for all $\alpha \in UB[g(b)]_{\cong_A} \cap S$; on the other hand, since $g(b) \in S$ then $g(b) \in p\text{-max}[g(b)]_{\cong_A}$, particularly $g(b) \in UB[g(b)]_{\cong_A}$, hence $g(b) \in UB[g(b)]_{\cong_A} \cap S$. As a result, we obtain $\rho_A(y, g(b)) = \top$. Now, connecting expression (7) with transitivity of ρ_A ,

$$\rho_A(a, g(b)) \leq \rho_A(x, y) = \rho_A(x, y) \otimes \rho_A(y, g(b)) \quad (8)$$

$$\leq \rho_A(x, g(b)) \quad (9)$$

for all $x \in \varphi_S(a)$. Joining Equations (6) and (8) we obtain, $\rho_A(x, g(b)) = \rho_A(a, g(b))$ and, finally, Equation (5) leads to

$$\rho_B^{a_0}(f(a), b) = \rho_A(a, g(b)). \quad \square$$

We can now conclude this section by stating the necessary and sufficient conditions for the existence of right adjoint from a fuzzy preorder to an unstructured set. In this statement, for readability reasons, we do not use the syntactic sugared version of the previous lemma (namely, φ_S) but, instead, state the conditions directly in their low level appearance.

Theorem 5: Given a fuzzy preordered set $\mathbb{A} = (A, \rho_A)$ together with a mapping $f: A \rightarrow B$, there exists a fuzzy preordering ρ_B in B and a mapping $g: B \rightarrow A$ such that $(f, g): \mathbb{A} \rightleftharpoons \mathbb{B}$ if and only if there exists $S \subseteq A$ such that, for all $a, a_1, a_2, \in A$:

- 1) $S \subseteq \bigcup_{a \in A} p\text{-max}[a]_{\cong_A}$
- 2) $p\text{-min}(UB[a]_{\cong_A} \cap S) \neq \emptyset$
- 3) $\rho_A(a_1, a_2) \leq (p\text{-min}(UB[a_1]_{\cong_A} \cap S) \sqsubseteq_H p\text{-min}(UB[a_2]_{\cong_A} \cap S))$.

PROOF: Necessity follows from [25, Thm. 4], considering $S = gf(A)$; sufficiency follows from Lemma 4. \square

V. CONCLUSIONS

Based on the set of necessary conditions for the existence of right adjoint (between fuzzy preorders) to a mapping $f: (A, \rho_A) \rightarrow B$, we have proved that these conditions are also sufficient.

It is remarkable the fact that the right adjoint is not unique. In fact, there is a number of degrees of freedom in order to define it: just consider the parameterized construction of g that we have given in terms of an element $a_0 \in A$ (in the case of non-surjective f). Note, however, that our results do not imply that every right adjoint should be like that; we simply chose a convenient construction to extent the induced fuzzy ordering on the image of f to the whole set B , and maybe other constructions would be adequate as well (but this is further work).

It is worth to note that there are different versions of antisymmetry and reflexivity in a fuzzy environment (see, for instance, [5], [7]). Accordingly, another line of future work will be the adaptation of the current results to these alternative definitions. Another source of future work will be to study the potential relationship to other approaches based on adequate versions of fuzzy closure systems [29].

ACKNOWLEDGMENT

Supported by Grants TIN2011-28084, TIN12-39353-C04-01, and TIN2014-59471-Pof the Science and Innovation Ministry of Spain, co-funded by the European Regional Development Fund (ERDF).

REFERENCES

- [1] R. Bělohlávek. Fuzzy Galois connections. *Mathematical Logic Quarterly*, 45(4):497–504, 1999.
- [2] R. Bělohlávek. Lattices of fixed points of fuzzy Galois connections. *Mathematical Logic Quarterly*, 47(1):111–116, 2001.
- [3] R. Bělohlávek. and P. Osíčka. Triadic fuzzy Galois connections as ordinary connections. In *IEEE Intl Conf on Fuzzy Systems*, 2012.
- [4] U. Bodenhofer. A similarity-based generalization of fuzzy orderings preserving the classical axioms. *International Journal of Uncertainty, Fuzziness and Knowledge-Based Systems* 8(5):593–610, 2000.
- [5] U. Bodenhofer. Representations and constructions of similarity-based fuzzy orderings. *Fuzzy Sets and Systems* 137(1):113–136, 2003.
- [6] U. Bodenhofer and F. Klawonn. A formal study of linearity axioms for fuzzy orderings. *Fuzzy Sets and Systems* 145(3):323–354, 2004.
- [7] U. Bodenhofer, B. De Baets and J. Fodor. A compendium of fuzzy weak orders: Representations and constructions. *Fuzzy Sets and Systems* 158(8):811–829, 2007.
- [8] F. Börner. Basics of Galois connections. *Lect. Notes in Computer Science*, 5250:38–67, 2008.
- [9] G. Castellini, J. Kosłowski, and G. E. Strecker. Categorical closure operators via Galois connections. *Mathematical research*, 67:72–72, 1992.
- [10] D. Cohen, P. Creed, P. Jeavons, and S. Živný. An algebraic theory of complexity for valued constraints: Establishing a Galois connection. *Lect. Notes in Computer Science*, 6907:231–242, 2011.
- [11] Cordero, P., Enciso, M., Mora, A., de Guzmán, I.P.: SLFD logic: Elimination of data redundancy in knowledge representation. *Lecture Notes in Computer Science* 2527 (2002) 141–150
- [12] Z. Csajbók and T. Mihálydeák. Partial approximative set theory: generalization of the rough set theory. *Intl J of Computer Information Systems and Industrial Management Applications*, 4:437–444, 2012.
- [13] B. Davey and H. Priestley. *Introduction to Lattices and Order*. Cambridge University Press, second edition, 2002.
- [14] K. Denecke, M. Erné, and S. L. Wismath. *Galois connections and applications*, volume 565. Springer, 2004.
- [15] R. P. Dilworth and M. Ward. Residuated lattices. *Transactions of the American Mathematical Society*, 45:335–354, 1939.
- [16] Y. Djouadi. Extended Galois derivation operators for information retrieval based on fuzzy formal concept lattice. *Lect. Notes in Computer Science*, 6929:346–358, 2011.
- [17] Y. Djouadi and H. Prade. Interval-valued fuzzy Galois connections: Algebraic requirements and concept lattice construction. *Fundamenta Informaticae*, 99(2):169–186, 2010.
- [18] W. Dzik, J. Järvinen, and M. Kondo. Intuitionistic propositional logic with Galois connections. *Logic Journal of the IGPL*, 18(6):837–858, 2010.
- [19] P. Filipiuk, M. Terepeta, H. Nielson, and F. Nielson. Galois connections for flow algebras. *Lect. Notes in Computer Science*, 6722:138–152, 2011.
- [20] J. Fodor and M. Roubens. *Fuzzy Preference Modelling and Multicriteria Decision Support*. Kluwer Academic Publishers, Dordrecht, 1994.
- [21] A. Frasca. Fuzzy Galois connections under weak conditions. *Fuzzy Sets and Systems*, 172(1):33–50, 2011.
- [22] L. Freeman and D. White. Using Galois lattices to represent network data. *Sociological Methodology*, pages 127–146, 1993.
- [23] J. G. García, I. Mardones-Pérez, M. A. de Prada-Vicente, and D. Zhang. Fuzzy Galois connections categorically. *Math. Log. Q.*, 56(2):131–147, 2010.
- [24] F. García-Pardo, I.P. Cabrera, P. Cordero, and M. Ojeda-Aciego. On Galois connections and Soft Computing. *Lect. Notes in Computer Science*, 7903:224–235, 2013.
- [25] F. García-Pardo, I.P. Cabrera, P. Cordero, and M. Ojeda-Aciego. On adjunctions between fuzzy preordered sets: necessary conditions. In *Proc. of Rough Sets and Current Trends in Computing*, 2014. To appear.
- [26] F. García-Pardo, I.P. Cabrera, P. Cordero, and M. Ojeda-Aciego. On the construction of fuzzy Galois connections. *Proc. of XVII Spanish Conference on Fuzzy Logic and Technology*, pages 99–102, 2014.
- [27] F. García-Pardo, I.P. Cabrera, P. Cordero, M. Ojeda-Aciego, and F.J. Rodríguez. Generating isotone Galois connections on an unstructured codomain. *Proc. of Information Processing and Management of Uncertainty in Knowledge-based Systems (IPMU)*, 2014. To appear.
- [28] F. García-Pardo, I.P. Cabrera, P. Cordero, M. Ojeda-Aciego, and F.J. Rodríguez. Building isotone Galois connections between preorders on an unstructured codomain. *Lect. Notes in Computer Science* 8478:67–79, 2014.
- [29] L. Guo, G.-Q. Zhang, and Q. Li. Fuzzy closure systems on L -ordered sets. *Mathematical Logic Quarterly*, 57(3):281–291, 2011.