

OBSERVATIONS OF THE 2008 SUMMER WILDFIRES BY GASP, MODIS, AND CALIPSO

Jasper R. Lewis

Center for Atmospheric Sciences, Hampton, VA 23668, USA

1. INTRODUCTION

Between June – August of 2008, northern California experienced a large number of wildfires dubbed the Northern California Lightning Series by the California Department of Forestry and Fire Protection. A severe thunderstorm, which produced over 6,000 lightning strikes, ignited 2,096 fires which burned nearly 1.2 million acres (4,856 km²). Extremely dry conditions caused by the first statewide drought in over a decade exasperated the problem, resulting in one of the worst wildfire situations in state history. Several exceedances of the National Ambient Air Quality Standards (NAAQS) for PM_{2.5} throughout the California Central Valley have been attributed to the wildfires. Such exceedances are a rarity during the summer within the region. Satellite measurements from the Moderate Resolution Imaging Spectroradiometer (MODIS) instrument aboard the Aqua satellite, GOES Aerosol/Smoke Product (GASP) from GOES-West, and Cloud-Aerosol Lidar with Orthogonal Polarization (CALIOP) aboard CALIPSO were used to show the impact on air quality from the wildfires.

2. METHODS AND RESULTS

Twenty-four-hour in-situ mass concentrations from eight surface PM_{2.5} monitors in the Sacramento Valley Air Basin (Chico, Yuba City, Sacramento, and UC Davis) and San Joaquin Valley Air Basin (Stockton, Fresno, Corcoran, and Bakersfield) were correlated with coincident aerosol optical depth (AOD) measurements from GASP and MODIS. Coincidences were defined as the nearest MODIS/GASP pixel within ± 0.05 -degrees latitude/longitude of the PM_{2.5} monitor. Correlations of PM_{2.5} and AOD in the western United States are typically poor due to high surface reflectances, variations in aerosol type, and higher occurrences of smoke and elevated aerosol plumes. In this analysis, the R²-correlations using MODIS and GASP were 0.47 and 0.23, respectively. Though




Figure 1. Correlations of PM_{2.5} to MODIS and GASP AOD at Chico (top) and Sacramento (bottom).

MODIS had stronger correlations than GASP, both instruments gave similar estimates of PM_{2.5} when a linear-fit model was used. Examples of the analysis for Chico and Sacramento are shown in Figure 1.

Finally, the vertical aerosol distributions determined from CALIOP were used identify elevated smoke plumes associated with transport. The analysis provides evidence that the PM_{2.5}-AOD relationship is improved when aerosol loading is contained primarily within the boundary layer.

3. SUMMARY

This presentation will demonstrate the ability of satellite instruments to monitor air quality during a large pollution event. The comparison of GASP, MODIS, and CALIPSO will highlight advantages of each instrument. The PM_{2.5}-AOD correlations found in this study varied by location, but was generally better in the Sacramento Valley Air Basin, which was nearest the origin of the wildfires, than in the San Joaquin Valley, where the majority of the smoke pollution came from transport. Stronger correlations were found using MODIS AOD than those found using GASP AOD. Typical values for the PM_{2.5}-AOD relationship were approximately $PM_{2.5} = 30 \tau + 9$ ($\mu\text{g m}^{-3}$), using a linear regression.